slender, an inch or two in length, covered with glandular hairs and supported by lanceolate bractes. Calyx divided into five ovate, concave, hairy segments ending in an acuminate point or a lanceolate leafet. Petals five, white, ovate or oblong, concave, contracted into a short claw at base. Stamens very numerous, with roundish anthers and slender, white filaments. Germs numerous, covering a conic central receptacle. Styles capillary, arising from the sides of the germs, persistent. Fruit a black, shining, compound berry formed of pulpy acini attached to the receptacle, each containing a single oblong seed.

The bark of the root of this bramble is the part which has been medicinally employed. It is a pure and strong astringent, which property it manifests both by its sensible effects and by chemical examination. When treated with the sulphate of iron both the tincture and decoction assume a beautiful dark purple colour and throw down a copious precipitate. A precipitate also takes place on the addition of gelatin, which is copious, white and opaque. The alcoholic solution is in part decomposed by water. The substance precipitated does not occasion the uniform turbidness which usually attends the separation

of resins, but exhibits a flocculent appearance like that of congulated mucilage. These flocculi, however, when collected and dried, exhibit the common resinous properties on exposure to heat. I subjected the dried bark to distillation, but the distilled water was nearly insipid, possessing only a very slight flavour of the root.

The properties of this bark are those of a very powerful astringent. I have tried its operation sufficiently to become satisfied of its efficacy both internally and externally used in a variety of cases which admit of relief from medicines of its class. It is true that our list of vegetable astringents has become very numerous and the cases which require them are perhaps less frequent than was formerly imagined; yet as we continue to import and consume various foreign medicines of this kind, we ought not to exclude from attention native articles of equal efficacy. Professor Chapman, of Philadelphia, expresses a very decided opinion in regard to the powers of this substance. "Of the vegetable astringents," says he, "this I have reason to believe is among the most active and decidedly efficacious in certain cases. To the declining stages of dysentery after the symptoms of active inflammation are removed, it is well suited, though I have given it, I think, with